



Dr. Fahri Yildiz,
ENT Specialist,
PAN Clinic, Zeppelinstr.1,
50667 Cologne, Germany

Anesthesia and Medication: 麻酔と薬剤:

One tablet (50 mg) Diclophenac was administered as peri-operative pain reduction and decongestion. If a patient had a strong gag reflex Lidocaine spray or Benzocaine gel was applied on the soft palate for the reduction of the gag reflex. For the local anesthesia of the soft palate, 0.5 ml Lidocaine 2 % with Adrenalin was injected into each of the positions (A) directly beneath the border between hard and soft palate (see Fig. 1). After 3 to 5 minutes, 0.5 ml was injected in each of the positions (B) above the palatal arcs (see Fig. 1). If required by the patient, for post-operative pain reduction and decongestion, 1-2 tablets (50 mg) Diclophenac during the day and 1 suppository (100 mg) Diclophenac during the night was administered for 1-2 days.

ジクロフェナク1錠(50mg)が周術期の疼痛抑制用と充血除去用に投与された。患者に強い絞扼反射がある場合は、リドカインスプレーまたはベンゾカインジェルが軟口蓋に投与された。軟口蓋の局部麻酔には、アドレナリン入りのリドカイン2% 0.5mlを硬口蓋と軟口蓋の境界の真下の各位置(A)に注射された(図1参照)。3~5分後、0.5mlが口蓋弓の上の各位置(B)に注射された(図1参照)。患者によっては、術後疼痛抑制と充血除去用に、日中にジクロフェナク1~2錠(50mg)と夜間に座薬1錠(100mg)が1~2日間投与された。

Fig.1 図1

Injections of anesthetics 麻酔剤の注射箇所




Fig.2 図2

Application of the probe フロープの適用箇所

